Excision in K-theory
Talk at the Institut Mittag-Leffler
April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

April 11, 2006

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Brave New Algebraic Geometry with singularities?

If we are going to embark on a world of Brave New Algebraic Geometry, [BNAG] we have to understand is local-to-global principles

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Brave New Algebraic Geometry with singularities?

If we are going to embark on a world of Brave New Algebraic Geometry, [BNAG] we have to understand is local-to-global principles

- **How do we assemble local information?**

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006
Bjørn Ian Dundas (joint with Harald Kittang)
Brave New Algebraic Geometry with singularities?
Motivating examples
Classical breakdown of excision
Damage control
Resolving the classical case
Extending to connective S-algebras
Questions and further work

Brave New Algebraic Geometry with singularities?

If we are going to embark on a world of Brave New Algebraic Geometry, [BNAG] we have to understand is local-to-global principles

- **How do we assemble local information?**
- **In particular: how do we paste vector bundles? (whose geometric interpretation in BNAG is still up for grabs)**

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Brave New Algebraic Geometry with singularities?

If we are going to embark on a world of Brave New Algebraic Geometry, [BNAG] we have to understand is local-to-global principles

- **How do we assemble local information?**
- **In particular: how do we paste vector bundles? (whose geometric interpretation in BNAG is still up for grabs)**

We even want to consider what can be said while allowing singularities [and so the questions are not accessible through motivic methods even in OAG].

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Topological K-theory

Topological K-theory is a cohomology theory and satisfies excision. If

$$\begin{array}{ccc} X_{12} & \longrightarrow & X_1 \\ \downarrow & & \downarrow \\ X_2 & \longrightarrow & X_0 \end{array}$$

is a (homotopy) cocartesian square of spaces, then

$$\begin{array}{ccc} K(X_{12}) & \longleftarrow & K(X_1) \\ \uparrow & & \uparrow \\ K(X_2) & \longleftarrow & K(X_0) \end{array}$$

is a homotopy cartesian square of spectra, illustrating that virtual bundles can be assembled from local data.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Topological K-theory

Topological K-theory is a cohomology theory and satisfies excision. If

$$\begin{array}{ccc} X_{12} & \longrightarrow & X_1 \\ \downarrow & & \downarrow \\ X_2 & \longrightarrow & X_0 \end{array}$$

is a (homotopy) cocartesian square of spaces, then

$$\begin{array}{ccc} K(X_{12}) & \longleftarrow & K(X_1) \\ \uparrow & & \uparrow \\ K(X_2) & \longleftarrow & K(X_0) \end{array}$$

is a homotopy cartesian square of spectra, illustrating that virtual bundles can be assembled from local data.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Algebraic K-theory

In AG one would consider a pushout square like

$$\begin{array}{ccc} X_{12} & \longrightarrow & X_1 \\ \downarrow & & \downarrow \\ X_2 & \longrightarrow & X_0 \end{array}$$

in schemes, and ask whether the corresponding square of algebraic K-theory spectra were homotopy cartesian.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Let us look at the affine case. Let

$$\mathcal{A} = \left\{ \begin{array}{ccc} A_0 & \longrightarrow & A_1 \\ \downarrow & & \downarrow \\ A_2 & \longrightarrow & A_{12} \end{array} \right\}$$

be a cartesian square of rings (corresponding in the commutative case to a cocartesian square

$$Spec(\mathcal{A}) = \left\{ \begin{array}{ccc} Spec(A_0) & \longleftarrow & Spec(A_1) \\ \uparrow & & \uparrow \\ Spec(A_2) & \longleftarrow & Spec(A_{12}) \end{array} \right\}$$

of schemes).
To assure homotopy (co)cartesianness assume all arrows in $\mathcal{A}$ surjective.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Let us look at the affine case. Let

$$\mathcal{A} = \left\{ \begin{array}{ccc} A_0 & \longrightarrow & A_1 \\ \downarrow & & \downarrow \\ A_2 & \longrightarrow & A_{12} \end{array} \right\}$$

be a cartesian square of rings (corresponding in the commutative case to a cocartesian square

$$Spec(\mathcal{A}) = \left\{ \begin{array}{ccc} Spec(A_0) & \longleftarrow & Spec(A_1) \\ \uparrow & & \uparrow \\ Spec(A_2) & \longleftarrow & Spec(A_{12}) \end{array} \right\}$$

of schemes).

To assure homotopy (co)cartesianness assume all arrows in $\mathcal{A}$ surjective.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Let us look at the affine case. Let

$$\mathcal{A} = \left\{ \begin{array}{ccc} A_0 & \longrightarrow & A_1 \\ \downarrow & & \downarrow \\ A_2 & \longrightarrow & A_{12} \end{array} \right\}$$

be a cartesian square of rings (corresponding in the commutative case to a cocartesian square

$$Spec(\mathcal{A}) = \left\{ \begin{array}{ccc} Spec(A_0) & \longleftarrow & Spec(A_1) \\ \uparrow & & \uparrow \\ Spec(A_2) & \longleftarrow & Spec(A_{12}) \end{array} \right\}$$

of schemes).
To assure homotopy (co)cartesianness assume all arrows in $\mathcal{A}$ surjective.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Let us look at the affine case. Let

$$\mathcal{A} = \left\{ \begin{array}{ccc} A_0 & \longrightarrow & A_1 \\ \downarrow & & \downarrow \\ A_2 & \longrightarrow & A_{12} \end{array} \right\}$$

be a cartesian square of rings (corresponding in the commutative case to a cocartesian square

$$Spec(\mathcal{A}) = \left\{ \begin{array}{ccc} Spec(A_0) & \longleftarrow & Spec(A_1) \\ \uparrow & & \uparrow \\ Spec(A_2) & \longleftarrow & Spec(A_{12}) \end{array} \right\}$$

of schemes).
To assure homotopy (co)cartesianness assume all arrows in $\mathcal{A}$ surjective.
do 4 examples

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Examples

- **The disjoint union:**

$$\begin{array}{ccc} A_1 \times A_2 & \longrightarrow & A_1 \\ \downarrow & & \downarrow \\ A_2 & \longrightarrow & 0 \end{array}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Examples

- **The disjoint union:**

$$\begin{array}{ccc} A_1 \times A_2 & \longrightarrow & A_1 \\ \downarrow & & \downarrow \\ A_2 & \longrightarrow & 0 \end{array}$$

- **The one-point union**

$$\begin{array}{ccc} k[x,y]/xy & \longrightarrow & k[x] \\ \downarrow & & \downarrow \\ k[y] & \longrightarrow & k \end{array}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work


Examples

- **The disjoint union:**

$$\begin{array}{ccc} A_1 \times A_2 & \longrightarrow & A_1 \\ \downarrow & & \downarrow \\ A_2 & \longrightarrow & 0 \end{array}$$

- **The one-point union**

$$\begin{array}{ccc} k[x,y]/xy & \longrightarrow & k[x] \\ \downarrow & & \downarrow \\ k[y] & \longrightarrow & k \end{array}$$



Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Examples (cont.)

- **The cusp:**

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow{\substack{x\mapsto t^3 \\ y\mapsto t^2}} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

- Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case


Extending to connective S-algebras

Questions and further work

Examples (cont.)

- **The cusp:**

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow{\substack{x\mapsto t^3 \\ y\mapsto t^2}} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case


Extending to connective S-algebras

Questions and further work

Examples (cont.)

- **The cusp:**

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow[y\mapsto t^2]{x\mapsto t^3} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



- **The Rim square:**

$$\begin{array}{ccc} \mathbf{Z}[C_p] & \longrightarrow & \mathbf{Z}[\zeta_p] \\ \downarrow & & \downarrow \\ \mathbf{Z} & \longrightarrow & \mathbf{F}_p \end{array}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Bass' K_1 and Milnor's theorem

Bass' program for higher K-theory rested on getting this right: he realized that the Whitehead groups K_1 could be built out of "clutching functions" so that excision seemed secure.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Bass' K_1 and Milnor's theorem

Bass' program for higher K-theory rested on getting this right: he realized that the Whitehead groups K_1 could be built out of "clutching functions" so that excision seemed secure.
at the heart of the matter: Milnor's theorem

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Bass' K_1 and Milnor's theorem

Bass' program for higher K-theory rested on getting this right: he realized that the Whitehead groups K_1 could be built out of "clutching functions" so that excision seemed secure.

Theorem [Milnor]

Let $\mathcal{A}$ be a cartesian square of rings where the vertical maps are surjections. Then the corresponding square of categories of finitely generated modules $\mathcal{P}_{\mathcal{A}}$ is a fiber square.

Excision in low dimensions

In the end Milnor establishes that if all maps in the cartesian square are surjections you get a Mayer-Vietoris sequence

$$K_2(A_1) \oplus K_2(A_2) \to K_2(A_{12}) \to$$
$$K_1(A_0) \to K_1(A_1) \oplus K_1(A_2) \to K_1(A_{12})$$
$$\to K_0(A_0) \to K_0(A_1) \oplus K_0(A_2)$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Excision in low dimensions

In the end Milnor establishes that if all maps in the cartesian square are surjections you get a Mayer-Vietoris sequence

$$K_2(A_1) \oplus K_2(A_2) \to K_2(A_{12}) \to K_1(A_0) \to K_1(A_1) \oplus K_1(A_2) \to K_1(A_{12}) \to K_0(A_0) \to K_0(A_1) \oplus K_0(A_2)$$

(without the K_2 and with half of the surjectivity this is in Bass).

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

The projective line

Non affine example:

$$\begin{array}{ccc} \mathcal{P}_{\mathbf{P}^1_k} & \longrightarrow & \mathcal{P}_{k[t]} \\ \downarrow & & \downarrow \\ \mathcal{P}_{k[t^{-1}]} & \longrightarrow & \mathcal{P}_{k[t,t^{-1}]} \end{array}$$

is a fiber square

Fact:

$$\begin{array}{ccc} K(\mathbf{P}^1_k) & \longrightarrow & K(k[t]) \\ \downarrow & & \downarrow \\ K(k[t^{-1}]) & \longrightarrow & K(k[t,t^{-1}]) \end{array}$$

is homotopy cartesian.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control


Resolving the classical case

Extending to connective S-algebras

Questions and further work

The projective line

$$\begin{array}{ccc} \mathcal{P}_{\mathbf{P}^1_k} & \longrightarrow & \mathcal{P}_{k[t]} \\ \downarrow & & \downarrow \\ \mathcal{P}_{k[t^{-1}]} & \longrightarrow & \mathcal{P}_{k[t,t^{-1}]} \end{array}$$



is a fiber square (the vector bundles on $\mathbf{P}^1_k$ are gotten by gluing together two bundles over affine lines).

Fact:

$$\begin{array}{ccc} K(\mathbf{P}^1_k) & \longrightarrow & K(k[t]) \\ \downarrow & & \downarrow \\ K(k[t^{-1}]) & \longrightarrow & K(k[t,t^{-1}]) \end{array}$$

is homotopy cartesian.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control


Resolving the classical case

Extending to connective S-algebras

Questions and further work

The projective line

$$\begin{array}{ccc} \mathcal{P}_{\mathbf{P}^1_k} & \longrightarrow & \mathcal{P}_{k[t]} \\ \downarrow & & \downarrow \\ \mathcal{P}_{k[t^{-1}]} & \longrightarrow & \mathcal{P}_{k[t,t^{-1}]} \end{array}$$



is a fiber square

Fact:

$$\begin{array}{ccc} K(\mathbf{P}^1_k) & \longrightarrow & K(k[t]) \\ \downarrow & & \downarrow \\ K(k[t^{-1}]) & \longrightarrow & K(k[t,t^{-1}]) \end{array}$$

is homotopy cartesian.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Classical breakdown of excision

However, Swan showed that there was no way this could continue: there is no functor K_3 extending this sequence:

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Example: the coordinate axes

$$\begin{array}{ccc} k[x,y]/xy & \longrightarrow & k[x] \\ \downarrow & & \downarrow \\ k[y] & \longrightarrow & k \end{array}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control


Resolving the classical case

Extending to connective S-algebras

Questions and further work

Example: the coordinate axes

$$\begin{array}{ccc} k[x,y]/xy & \longrightarrow & k[x] \\ \downarrow & & \downarrow \\ k[y] & \longrightarrow & k \end{array}$$



Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control


Resolving the classical case

Extending to connective S-algebras

Questions and further work

Example: the coordinate axes

$$\begin{array}{ccc} k[x,y]/xy & \longrightarrow & k[x] \\ \downarrow & & \downarrow \\ k[y] & \longrightarrow & k \end{array}$$



With $k = \mathbf{Z}$ you get that $K_2 k[x,y]/xy \cong K_2 k \oplus \mathbf{Z}$.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control


Resolving the classical case

Extending to connective S-algebras

Questions and further work

Example: the coordinate axes

$$\begin{array}{ccc} k[x,y]/xy & \longrightarrow & k[x] \\ \downarrow & & \downarrow \\ k[y] & \longrightarrow & k \end{array}$$



With $k = \mathbf{Z}$ you get that $K_2 k[x,y]/xy \cong K_2 k \oplus \mathbf{Z}$ (where the last factor is generated by $[x_{12}^x, x_{21}^y] \in K_2 = ker\{St \to E\}$ if I remember correctly).

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control


Resolving the classical case

Extending to connective S-algebras

Questions and further work

Example: the coordinate axes

$$\begin{array}{ccc} k[x,y]/xy & \longrightarrow & k[x] \\ \downarrow & & \downarrow \\ k[y] & \longrightarrow & k \end{array}$$



With $k = \mathbf{Z}$ you get that $K_2 k[x,y]/xy \cong K_2 k \oplus \mathbf{Z}$. But then the fact that the surjections are split and functoriality of K_3 forces the boundary map in

$$K_3(k[x]) \oplus K_3(k[y]) \to K_3(k) \to K_2(k[x,y]/xy) \to K_2(k[x]) \oplus K_2(k[y])$$

to be zero,

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control


Resolving the classical case

Extending to connective S-algebras

Questions and further work

Example: the coordinate axes

$$\begin{array}{ccc} k[x,y]/xy & \longrightarrow & k[x] \\ \downarrow & & \downarrow \\ k[y] & \longrightarrow & k \end{array}$$



With $k = \mathbf{Z}$ you get that $K_2 k[x,y]/xy \cong K_2 k \oplus \mathbf{Z}$. But then the fact that the surjections are split and functoriality of K_3 forces the boundary map in

$$K_3(k[x]) \oplus K_3(k[y]) \to K_3(k) \to K_2(k[x,y]/xy) \to K_2(k[x]) \oplus K_2(k[y])$$

to be zero, but the infinite cyclic factor of $K_2(k[x,y]/xy)$ can't map injectively into $K_2(k[x]) \oplus K_2(k[y]) \cong \mathbf{Z}/2 \oplus \mathbf{Z}/2$.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Damage control

So, Milnor's theorem is not enough: that the square of exact categories is a fiber square does not guarantee that its K-theory will be homotopy cartesian.
Can one control this breakdown of excision?

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006
Bjørn Ian Dundas (joint with Harald Kittang)
Brave New Algebraic Geometry with singularities?
Motivating examples
Classical breakdown of excision
Damage control
Resolving the classical case
Extending to connective S-algebras
Questions and further work

First step in this direction came around 1980 when Guin-Waléry and Loday showed (with a slight reinterpretation) that if

$$\mathcal{A} = \left\{ \begin{array}{ccc} A_0 & \longrightarrow & A_1 \\ \downarrow & & \downarrow f_1 \\ A_2 & \xrightarrow{f_2} & A_{12} \end{array} \right\}$$

is a cartesian square of Q-algebras with all maps surjective and $I_j = ker\{f_j\}$, then the first nonvanishing obstruction group satisfies

$$\pi_2 ifiK(\mathcal{A}) \cong I_1/I_1^2 \otimes_{A_0^o \otimes A_0} I_2/I_2^2 \cong \pi_1 ifiHC(\mathcal{A})$$

where *ifi* is the iterated fiber of a cube of spectra, and *HC* is cyclic homology.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

The KABI-conjecture

It was soon conjectured (at least by Weibel, and perhaps by others) that if A_0 is a Q-algebra, then

$$\pi_q ifiK(\mathcal{A}) \cong \pi_{q-1} ifiHC(A)$$

for all q .

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

The KABI-conjecture

It was soon conjectured (at least by Weibel, and perhaps by others) that if A_0 is a Q-algebra, then

$$\pi_q ifiK(\mathcal{A}) \cong \pi_{q-1} ifiHC(A)$$

for all q [give a brief history of the KABI-conjecture].

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Resolving the classical case

The conjecture remained unresolved until recently when Cortiñas proved it using the techniques of Suslin and Wodzicki [Ann. Math. 92].

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006
Bjørn Ian Dundas (joint with Harald Kittang)
Brave New Algebraic Geometry with singularities?
Motivating examples
Classical breakdown of excision
Damage control
Resolving the classical case
Extending to connective S-algebras
Questions and further work

Resolving the classical case

Theorem [Cortiñas, Inv. math. 06]

If $\mathcal{A}$ is a cartesian square of Q-algebras with vertical maps surjective, then the trace map induces an equivalence of spectra

$$ifiK(\mathcal{A}) \xrightarrow{\sim} ifiHC^{-}(\mathcal{A}) \xleftarrow{\sim} S^1 \wedge ifiHC(\mathcal{A}).$$

Considering a map of squares as a cube, Cortiñas theorem says that both the cubes

$$K(\mathcal{A}) \to HC^{-}(\mathcal{A}), \text{ and } S^1 \wedge HC(\mathcal{A}) \to HC^{-}(\mathcal{A})$$

are homotopy cartesian.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Resolving the classical case

Theorem [Cortiñas, Inv. math. 06]

If $\mathcal{A}$ is a cartesian square of Q-algebras with vertical maps surjective, then the trace map induces an equivalence of spectra

$$ifiK(\mathcal{A}) \xrightarrow{\sim} ifiHC^{-}(\mathcal{A}) \xleftarrow{\sim} S^1 \wedge ifiHC(\mathcal{A}).$$

Considering a map of squares as a cube, Cortiñas theorem says that both the cubes

$$K(\mathcal{A}) \to HC^{-}(\mathcal{A}), \text{ and } S^1 \wedge HC(\mathcal{A}) \to HC^{-}(\mathcal{A})$$

are homotopy cartesian.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Resolving the classical case

Theorem [Cortiñas, Inv. math. 06]

If $\mathcal{A}$ is a cartesian square of Q-algebras with vertical maps surjective, then the trace map induces an equivalence of spectra

$$ifiK(\mathcal{A}) \xrightarrow{\sim} ifiHC^{-}(\mathcal{A}) \xleftarrow{\sim} S^1 \wedge ifiHC(\mathcal{A}).$$

Considering a map of squares as a cube, Cortiñas theorem says that both the cubes

$$K(\mathcal{A}) \to HC^{-}(\mathcal{A}), \text{ and } S^1 \wedge HC(\mathcal{A}) \to HC^{-}(\mathcal{A})$$

are homotopy cartesian.
The connection between cyclic homology and negative cyclic homology derives from the fact that periodic cyclic homology does satisfy excision ($HP(\mathcal{A})$ is homotopy cartesian).

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Geller, Reid and Weibel studied the cyclic homology side of the story quite extensively in the 80's, and so this leads to immediate calculations:

Geller, Reid and Weibel studied the cyclic homology side of the story quite extensively in the 80's, and so this leads to immediate calculations:

Example, the cusp

Let k be a field of characteristic 0

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow[y\mapsto t^2]{x\mapsto t^3} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras


Questions and further work

Geller, Reid and Weibel studied the cyclic homology side of the story quite extensively in the 80's, and so this leads to immediate calculations:

Example, the cusp

Let k be a field of characteristic 0

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow[y\mapsto t^2]{x\mapsto t^3} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case


Extending to connective S-algebras

Questions and further work

Example, the cusp

Let k be a field of characteristic 0

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow[y\mapsto t^2]{x\mapsto t^3} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



The fiber of $Kk \to K(k[t]/t^2)$ is known in terms of HC, [Goodwillie Ann. Math. 85]

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case


Extending to connective S-algebras

Questions and further work

Example, the cusp

Let k be a field of characteristic 0

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow[y\mapsto t^2]{x\mapsto t^3} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



The fiber of $Kk \to K(k[t]/t^2)$ is known in terms of HC, and so modulo knowing Kk the calculation of the K-theory of $C = k[x,y]/x^2 - y^3$ is reduced to HC-calculations [GRW, Crelle 89]:

$$\begin{aligned} K_0 C &= \mathbf{Z} \oplus k \\ K_1 C &= k^* \oplus \Omega^1_{k/\mathbf{Q}} \\ K_2 C &= K_2 k \oplus k[t]/t^2 \oplus \Omega^2_{k/\mathbf{Q}} \\ K_3 C &= K_3 k \oplus k[t]/t^2 \otimes_k \Omega^1_{k/\mathbf{Q}} \oplus \Omega^3_{k/\mathbf{Q}} \end{aligned}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case


Extending to connective S-algebras

Questions and further work

Example, the cusp

Let k be a field of characteristic 0

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow[y\mapsto t^2]{x\mapsto t^3} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



The fiber of $Kk \to K(k[t]/t^2)$ is known in terms of HC, and so modulo knowing Kk the calculation of the K-theory of $C = k[x,y]/x^2 - y^3$ is reduced to HC-calculations [GRW, Crelle 89]:

$$\begin{aligned} K_0 C &= \mathbf{Z} \oplus k \\ K_1 C &= k^* \oplus \Omega^1_{k/\mathbf{Q}} \\ K_2 C &= K_2 k \oplus k[t]/t^2 \oplus \Omega^2_{k/\mathbf{Q}} \\ K_3 C &= K_3 k \oplus k[t]/t^2 \otimes_k \Omega^1_{k/\mathbf{Q}} \oplus \Omega^3_{k/\mathbf{Q}} \end{aligned}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case


Extending to connective S-algebras

Questions and further work

Example, the cusp

Let k be a field of characteristic 0

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow[y\mapsto t^2]{x\mapsto t^3} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



The fiber of $Kk \to K(k[t]/t^2)$ is known in terms of HC, and so modulo knowing Kk the calculation of the K-theory of $C = k[x,y]/x^2 - y^3$ is reduced to HC-calculations [GRW, Crelle 89]:

$$\begin{aligned} K_0 C &= \mathbf{Z} \oplus k \\ K_1 C &= k^* \oplus \Omega^1_{k/\mathbf{Q}} \\ K_2 C &= K_2 k \oplus k[t]/t^2 \oplus \Omega^2_{k/\mathbf{Q}} \\ K_3 C &= K_3 k \oplus k[t]/t^2 \otimes_k \Omega^1_{k/\mathbf{Q}} \oplus \Omega^3_{k/\mathbf{Q}} \end{aligned}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case


Extending to connective S-algebras

Questions and further work

Example, the cusp

Let k be a field of characteristic 0

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow[y\mapsto t^2]{x\mapsto t^3} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



The fiber of $Kk \to K(k[t]/t^2)$ is known in terms of HC, and so modulo knowing Kk the calculation of the K-theory of $C = k[x,y]/x^2 - y^3$ is reduced to HC-calculations [GRW, Crelle 89]:

$$\begin{aligned} K_0 C &= \mathbf{Z} \oplus k \\ K_1 C &= k^* \oplus \Omega^1_{k/\mathbf{Q}} \\ K_2 C &= K_2 k \oplus k[t]/t^2 \oplus \Omega^2_{k/\mathbf{Q}} \\ K_3 C &= K_3 k \oplus k[t]/t^2 \otimes_k \Omega^1_{k/\mathbf{Q}} \oplus \Omega^3_{k/\mathbf{Q}} \end{aligned}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case


Extending to connective S-algebras

Questions and further work

Example, the cusp

Let k be a field of characteristic 0

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow[y\mapsto t^2]{x\mapsto t^3} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



The fiber of $Kk \to K(k[t]/t^2)$ is known in terms of HC, and so modulo knowing Kk the calculation of the K-theory of $C = k[x,y]/x^2 - y^3$ is reduced to HC-calculations [GRW, Crelle 89]:

$$\begin{aligned} K_0 C &= \mathbf{Z} \oplus k \\ K_1 C &= k^* \oplus \Omega^1_{k/\mathbf{Q}} \\ K_2 C &= K_2 k \oplus k[t]/t^2 \oplus \Omega^2_{k/\mathbf{Q}} \\ K_3 C &= K_3 k \oplus k[t]/t^2 \otimes_k \Omega^1_{k/\mathbf{Q}} \oplus \Omega^3_{k/\mathbf{Q}} \end{aligned}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case


Extending to connective S-algebras

Questions and further work

Example, the cusp

Let k be a field of characteristic 0

$$\begin{array}{ccc} k[x,y]/x^2-y^3 & \xrightarrow{\substack{x\mapsto t^3\\ y\mapsto t^2}} & k[t] \\ \downarrow & & \downarrow \\ k & \longrightarrow & k[t]/t^2 \end{array}$$



The fiber of $Kk \to K(k[t]/t^2)$ is known in terms of HC, and so modulo knowing Kk the calculation of the K-theory of $C = k[x,y]/x^2 - y^3$ is reduced to HC-calculations [GRW, Crelle 89]:

$$K_n\mathbf{Q}[x,y]/x^2-y^3 = \begin{cases} \mathbf{Z}\oplus\mathbf{Q} & \text{if } n=0 \\ K_n(\mathbf{Q})\oplus\mathbf{Q}[t]/t^2 & \text{if } n=2j>0 \\ 0 & \text{otherwise} \end{cases}$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Positive characteristic

Outside characteristic zero, cyclic homology must of course be exchanged for topological cyclic homology TC**:**

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Positive characteristic

Outside characteristic zero, cyclic homology must of course be exchanged for topological cyclic homology TC:

Theorem [Geisser-Hesselholt, Inv. math. 06]

If $\mathcal{A}$ is a cartesian square of rings with vertical maps surjective, then the cyclotomic trace map induces a cube

$$K(\mathcal{A}) \to TC(\mathcal{A})$$

which is homotopy cartesian after profinite completion.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Positive characteristic

Outside characteristic zero, cyclic homology must of course be exchanged for topological cyclic homology TC:

Theorem [Geisser-Hesselholt, Inv. math. 06]

If $\mathcal{A}$ is a cartesian square of rings with vertical maps surjective, then the cyclotomic trace map induces a cube

$$K(\mathcal{A}) \to TC(\mathcal{A})$$

which is homotopy cartesian after profinite completion.

Since topological cyclic homology is open to attack with tools from stable homotopy theory, this makes calculations of K-theory possible.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Positive characteristic

Outside characteristic zero, cyclic homology must of course be exchanged for topological cyclic homology TC:

Theorem [Geisser-Hesselholt, Inv. math. 06]

If $\mathcal{A}$ is a cartesian square of rings with vertical maps surjective, then the cyclotomic trace map induces a cube

$$K(\mathcal{A}) \to TC(\mathcal{A})$$

which is homotopy cartesian after profinite completion.

Since topological cyclic homology is open to attack with tools from stable homotopy theory, this makes calculations of K-theory possible. For instance, Geisser and Hesselholt have calculated the K-theory of the coordinate axes $\mathbf{F}_p[x, y]/xy$.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Extending to connective **S**-algebras

Since simplicial rings lie densely in connective S-algebras, one would expect that these results followed for connective S-algebras once one extended the results from discrete rings to simplicial rings.

Indeed this is true, and such a proof has been worked out. However, there is a quicker proof, which I will present shortly.

Reason for wanting the long proof: Simplicial rings give hope of uncovering the right homotopical conditions on the categories of modules (surjectivity is not the right condition).

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Extending to connective **S**-algebras

Since simplicial rings lie densely in connective S-algebras, one would expect that these results followed for connective S-algebras once one extended the results from discrete rings to simplicial rings.
Indeed this is true, and such a proof has been worked out. However, there is a quicker proof, which I will present shortly.
Reason for wanting the long proof: Simplicial rings give hope of uncovering the right homotopical conditions on the categories of modules (surjectivity is not the right condition).

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006
Bjørn Ian Dundas (joint with Harald Kittang)
Brave New Algebraic Geometry with singularities?
Motivating examples
Classical breakdown of excision
Damage control
Resolving the classical case
Extending to connective S-algebras
Questions and further work

Extending to connective **S**-algebras

Since simplicial rings lie densely in connective S-algebras, one would expect that these results followed for connective S-algebras once one extended the results from discrete rings to simplicial rings.
Indeed this is true, and such a proof has been worked out. However, there is a quicker proof, which I will present shortly.
Reason for wanting the long proof: Simplicial rings give hope of uncovering the right homotopical conditions on the categories of modules (surjectivity is not the right condition).

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006
Bjørn Ian Dundas (joint with Harald Kittang)
Brave New Algebraic Geometry with singularities?
Motivating examples
Classical breakdown of excision
Damage control
Resolving the classical case
Extending to connective S-algebras
Questions and further work

Extending to connective **S**-algebras

Theorem [D, Kittang]

If $\mathcal{A}$ is a cartesian square of connective S-algebras with vertical maps 0-connected, then the cyclotomic trace map induces a homotopy cartesian cube

$$K(\mathcal{A}) \to TC(\mathcal{A}).$$

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Extending to connective **S**-algebras

Theorem [D, Kittang]

If $\mathcal{A}$ is a cartesian square of connective S-algebras with vertical maps 0-connected, then the cyclotomic trace map induces a homotopy cartesian cube

$$K(\mathcal{A}) \to TC(\mathcal{A}).$$

Note:

The circulating preprint only claims this after profinite completion, but using Goodwillie's model for integral TC and the techniques of Suslin/Wodzicki and Geisser/Hesselholt, Kittang has worked out the integral statement.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Proof

Let F be the homotopy fiber of the p-completion of the cyclotomic trace $K \to TC$.
The theorem is equivalent to the claim that $F(\mathcal{A})$ is homotopy cartesian, that is, F satisfies excision.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Proof

Let F be the homotopy fiber of the p-completion of the cyclotomic trace $K \to TC$.
The theorem is equivalent to the claim that $F(\mathcal{A})$ is homotopy cartesian
The result is easily proven from

Proof

Let F be the homotopy fiber of the p-completion of the cyclotomic trace $K \to TC$.
The theorem is equivalent to the claim that $F(\mathcal{A})$ is homotopy cartesian
The result is easily proven from

Lemma

Under the assumptions of the theorem,

$$\pi_0 A_0 \to \pi_0 A_1 \times_{\pi_0 A_{12}} \pi_0 A_2$$

is a surjection with square zero kernel.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Let F be the homotopy fiber of the p-completion of the cyclotomic trace $K \to TC$.

Lemma

Under the assumptions of the theorem,

$$\pi_0 A_0 \to \pi_0 A_1 \times_{\pi_0 A_{12}} \pi_0 A_2$$

is a surjection with square zero kernel.

McCarthy's theorem + lemma:

$$F(\pi_0 A_0) \to F(\pi_0 A_1 \times_{\pi_0 A_{12}} \pi_0 A_2)$$

is an equivalence

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Let F be the homotopy fiber of the p-completion of the cyclotomic trace $K \to TC$.

Lemma

Under the assumptions of the theorem,

$$\pi_0 A_0 \to \pi_0 A_1 \times_{\pi_0 A_{12}} \pi_0 A_2$$

is a surjection with square zero kernel.

McCarthy's theorem + lemma:

$$F(\pi_0 A_0) \to F(\pi_0 A_1 \times_{\pi_0 A_{12}} \pi_0 A_2)$$

is an equivalence

Assumptions and Geisser-Hesselholt:

$$\begin{array}{ccc} F(\pi_0 A_1 \times_{\pi_0 A_{12}} \pi_0 A_2) & \longrightarrow & F(\pi_0 A_1) \\ \downarrow & & \downarrow \\ F(\pi_0 A_2) & \longrightarrow & F(\pi_0 A_{12}) \end{array}$$

is homotopy cartesian

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

I.e.,

$F(\pi_0\mathcal{A})$ **is homotopy cartesian.**

Hence $F(\mathcal{A})$ is homotopy cartesian.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

I.e.,

$F(\pi_0\mathcal{A})$ **is homotopy cartesian.**

Denseness:

each node in the square of maps

$$F(\mathcal{A}) \to F(\pi_0(\mathcal{A}))$$

is an equivalence.

Hence $F(\mathcal{A})$ is homotopy cartesian.

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

I.e.,

$F(\pi_0\mathcal{A})$ **is homotopy cartesian.**

Denseness:

each node in the square of maps

$$F(\mathcal{A}) \to F(\pi_0(\mathcal{A}))$$

is an equivalence.

Hence $F(\mathcal{A})$ **is homotopy cartesian.**

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Questions

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Questions

- **What are the right conditions for the theorem?**

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006

Bjørn Ian Dundas (joint with Harald Kittang)

Brave New Algebraic Geometry with singularities?

Motivating examples

Classical breakdown of excision

Damage control

Resolving the classical case

Extending to connective S-algebras

Questions and further work

Questions

- **What are the right conditions for the theorem?**
- **What does the theorem really measure in terms of group completions?**

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006
Bjørn Ian Dundas (joint with Harald Kittang)
Brave New Algebraic Geometry with singularities?
Motivating examples
Classical breakdown of excision
Damage control
Resolving the classical case
Extending to connective S-algebras
Questions and further work

Questions

- **What are the right conditions for the theorem?**
- **What does the theorem really measure in terms of group completions?**
- **Does this have any implications for A-theory and manifolds?**

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006
Bjørn Ian Dundas (joint with Harald Kittang)
Brave New Algebraic Geometry with singularities?
Motivating examples
Classical breakdown of excision
Damage control
Resolving the classical case
Extending to connective S-algebras
Questions and further work

Questions

- What are the right conditions for the theorem?
- What does the theorem really measure in terms of group completions?
- Does this have any implications for A-theory and manifolds?
- What replaces the "Rim squares" in S-algebras?

Excision in K-theory Talk at the Institut Mittag-Leffler April 2006
Bjørn Ian Dundas (joint with Harald Kittang)
Brave New Algebraic Geometry with singularities?
Motivating examples
Classical breakdown of excision
Damage control
Resolving the classical case
Extending to connective S-algebras
Questions and further work

Questions

- What are the right conditions for the theorem?
- What does the theorem really measure in terms of group completions?
- Does this have any implications for A-theory and manifolds?
- What replaces the "Rim squares" in S-algebras?
- What does it tell us about vector bundles over S-algebras?

Questions

- What are the right conditions for the theorem?
- What does the theorem really measure in terms of group completions?
- Does this have any implications for A-theory and manifolds?
- What replaces the "Rim squares" in S-algebras?
- What does it tell us about vector bundles over S-algebras?
- Is a study of "combinatorial" S-algebras possible? (e.g., what are BN Stanley-Reisner rings"?)