

Profile of Two-Higgs-Doublet-Model Parameter Space

Abdul Wahab El Kaffas¹, Odd Magne OGREID² and Per Osland¹

1- Department of Physics and Technology, University of Bergen
Postboks 7803, N-5020 Bergen, Norway

2- Bergen University College, Bergen, Norway

We review recent work on constraining the parameter space of the Two-Higgs-Doublet Model by theoretical and experimental results. Some characteristics of the model, in particular the distribution of masses in the surviving parameter space, are discussed.

1 Introduction

We report on recent work on constraining the multi-dimensional parameter space of the Two-Higgs-Doublet Model by theoretical and experimental results [1, 2].

As compared with the Standard Model (SM), the Two-Higgs-Doublet Model (2HDM) allows for an additional mechanism for CP violation [3]. This is one of the main reasons for continued strong interest in the model [4].

Several experimental constraints restrict its parameter space. The $B \rightarrow X_s \gamma$ rate excludes low values of the charged-Higgs mass, $M_{H^\pm}$ [5], whereas $B - \bar{B}$ oscillations and the branching ratio R_b for $Z \rightarrow b\bar{b}$ exclude low values of $\tan\beta$. The precise measurements at LEP of the ρ parameter constrain the mass splitting in the Higgs sector, and force the masses to be not far from the Z mass scale [6].

From the theoretical point of view, there are also consistency conditions. The potential has to be positive for large values of the fields [7, 8]. Furthermore, we require the tree-level Higgs-Higgs scattering amplitudes to be unitary [9]. Together, these constraints dramatically reduce the allowed parameter space of the model. In particular, the unitarity constrain excludes large values of $\tan\beta$, *unless* μ is reasonably large. This limit is basically the decoupling limit [10].

Our recent study [2], restricted to the so-called “Type II” version, where up-type and down-type quarks couple to different Higgs doublets, uses rather complete and up-to-date experimental results, as well as accurate theoretical predictions for the above quantities. We consider a model with the Z_2 symmetry respected by the quartic couplings, i.e., no λ_6 and λ_7 couplings. Otherwise, we allow for full generality. In particular, we allow for CP violation, taking λ_5 complex. (For a definition of the potential, see [2].) The neutral Higgs boson sector will thus contain three bosons, described by a 3×3 mixing matrix R . These three neutral Higgs bosons will in general all have CP-violating Yukawa couplings. A related study, focused more on large values of $\tan\beta$, was also presented at this Workshop [11].

2 Results

We parametrize the model in terms of the masses of the two lightest neutral Higgs bosons, together with the charged Higgs boson mass, $\tan\beta$, the soft parameter μ^2 , and the rotation matrix R of the neutral sector. The third (heaviest) neutral mass is then calculable, as well as the quartic couplings, λ_i (see [12, 13]).

We establish allowed regions in the $\tan\beta$ - $M_{H^\pm}$ plane by the following procedure: For each point in this plane, we scan over the parameters $\alpha = \{\alpha_1, \alpha_2, \alpha_3\}$, defining the mixing matrix R in the neutral-Higgs sector, imposing the absolute theory constraints of positivity and unitarity. At each point, we evaluate a χ^2 penalty corresponding to the experimental constraints, adopting the “best” point (lowest χ^2) in α .

For two values of μ (200 and 500 GeV), we show in Fig. 1 the allowed regions in the $\tan\beta$ - $M_{H^\pm}$ plane, taking into account the theoretical constraints mentioned above, the LEP2 non-discovery, the very precise $\Delta\rho$ measurements at LEP, as well as the B -physics constraints ($B \rightarrow X_s \gamma$, mainly), and R_b . The masses of the two lightest neutral Higgs bosons are here kept fixed, at $M_1 = 100$ GeV and $M_2 = 300$ GeV or 500 GeV.

The over-all surviving regions of parameter space depend significantly on the “soft” parameter μ^2 . At low or negative values, the unitarity constraint will cut off the allowed region already at moderate values of $\tan\beta$. We have therefore shown results for a couple of positive values of μ^2 , the higher one approaching the so-called decoupling limit.

3 Distribution of Higgs masses

It turns out that, if μ is comparable with M_2 , or smaller, the distribution of M_3 -values will be very narrow, especially at large values of $\tan\beta$. This is illustrated in Fig. 2, for $M_1 = 100$ GeV, and two sets of (M_2, μ) values: (300, 200) GeV and (500, 500) GeV. Also, we note that for $M_2 = 500$ GeV and $\mu = 500$ GeV (lower panels), low values of $M_{H^\pm}$ are excluded. This is basically because of the $\Delta\rho$ constraint.




Figure 1: Allowed regions in the $\tan\beta$ - $M_{H^\pm}$ plane, taking into account theoretical and experimental constraints.




Figure 2: Distribution of M_3 -values for fixed $M_1 = 100$ GeV. Top: $M_2 = 300$ GeV and $\mu = 200$ GeV; bottom: $M_2 = 500$ GeV and $\mu = 500$ GeV. Three slices of $\tan\beta$ -values are shown.

On the other hand, if μ is larger than M_2 , the distribution can be considerably wider, as is seen in Fig. 3.

4 Summary

We have shown that the constraints of positivity and tree-level unitarity of Higgs–Higgs scattering, B -physics results, together with the precise LEP measurements, in particular of the ρ -parameter at LEP, exclude large regions of the 2HDM (II) parameter space. High values of $\tan\beta$ are excluded unless μ is large, allowing M_2 and M_3 both to be heavy. Furthermore, M_2 and M_3 should be reasonably close to each other. Improved precision of the $\bar{B} \rightarrow X_s \gamma$ measurement could significantly reduce the remaining part of the parameter space, but it appears unlikely that the model could be excluded other than by a negative search at the LHC.




Figure 3: Distribution of M_3 -values for $M_1 = 100$ GeV, $M_2 = 300$ GeV and $\mu = 500$ GeV. Three slices of $\tan \beta$ -values are shown, increasing to the right.

Acknowledgments

This research has been supported in part by the Mission Department of Egypt and the Research Council of Norway.

References

- [1] Slides:
<http://ilcagenda.linearcollider.org/contributionDisplay.py?contribId=151&sessionId=71&confId=1296>
- [2] A. W. El Kaffas, P. Osland and O. Magne OGREID, Phys. Rev. D, in print, arXiv:0706.2997 [hep-ph].
- [3] T. D. Lee, Phys. Rev. D **8**, 1226 (1973); S. Weinberg, Phys. Rev. Lett. **37**, 657 (1976).
- [4] E. Accomando *et al.*, “Workshop on CP studies and non-standard Higgs physics,” arXiv:hep-ph/0608079.
- [5] M. Misiak *et al.*, Phys. Rev. Lett. **98**, 022002 (2007) [arXiv:hep-ph/0609232].
- [6] S. Bertolini, Nucl. Phys. B **272**, 77 (1986).
- [7] N. G. Deshpande and E. Ma, Phys. Rev. D **18**, 2574 (1978); S. Nie and M. Sher, Phys. Lett. B **449**, 89 (1999) [arXiv:hep-ph/9811234]; S. Kanemura, T. Kasai and Y. Okada, Phys. Lett. B **471**, 182 (1999) [arXiv:hep-ph/9903289].
- [8] A. W. El Kaffas, W. Khater, O. M. OGREID and P. Osland, Nucl. Phys. B, **775**, 45 (2007) [arXiv:hep-ph/0605142].
- [9] S. Kanemura, T. Kubota and E. Takasugi, Phys. Lett. B **313**, 155 (1993) [arXiv:hep-ph/9303263]; A. G. Akeroyd, A. Arhrib and E. M. Naimi, Phys. Lett. B **490**, 119 (2000) [arXiv:hep-ph/0006035]; A. Arhrib, arXiv:hep-ph/0012353; I. F. Ginzburg and I. P. Ivanov, arXiv:hep-ph/0312374; Phys. Rev. D **72**, 115010 (2005) [arXiv:hep-ph/0508020].
- [10] J. F. Gunion and H. E. Haber, Phys. Rev. D **67**, 075019 (2003) [arXiv:hep-ph/0207010].
- [11] M. Krawczyk, in:
<http://ilcagenda.linearcollider.org/contributionDisplay.py?contribId=454&sessionId=71&confId=1296>
- [12] W. Khater and P. Osland, Nucl. Phys. B **661**, 209 (2003) [arXiv:hep-ph/0302004].
- [13] A. W. E. Kaffas, O. M. OGREID and P. Osland, Nonlinear Phenomena in Complex Systems, to appear [arXiv:hep-ph/0702097].